

A one-year comparison of new wind atlases over the North Sea

Etienne Cheynet, Ida Marie Solbrekke, Jan Markus Diezel, Joachim Reuder

Geophysical Institute and Bergen Offshore Wind Centre, University of Bergen



UNIVERSITETET I BERGEN

What is the presentation about?

(Climatological) Wind atlas:

Database containing wind speed and direction at specific height(s) over a long period (e.g. 30 years) in a given region.

Examples:

- ERA5 Reanalysis: 31 km horizontal resolution, 1-h time resolution, World
- NEWA: 3 km horizontal resolution, 30-min time resolution, Europe
- NORA3: 3 km horizontal resolution, 1-h time resolution, northern Europe




Figure: Hourly mean wind speed at 100 m above sea level (asl) from ERA5, between 2010 and 2020

Objective

To assess the performances of NORA3 and NEWA
in a case study (FINO1 in 2009)

Impact & applications

Wind Resource assessment

+

Wind turbine design

NORA3 & NEWA databases are open-access



Catalog <https://thredds.met.no/thredds/projects/nora3.html>

Terms of service

This is a shared public service which may experience overload in traffic from time to time. We reserve the right to block IP addresses if they cause traffic overload. Be nice: don't spawn multiple parallel downloads, our servers can handle a huge amount of traffic but may struggle if you start parallel downloads, especially if you don't wait for each request to answer before the next request is sent. If you require priority access for your own operational service please contact us at servicedesk@met.no. Get messages about planned maintenance and incidents from our [status page](#).

[MET Norway's Privacy Policy](#)

Dataset	Size	Last Modified
NORAS		--
NORAS/		--

[MET Norway Thredds Service at Norwegian Meteorological Institute](#) see [Info Documentation](#)



Contents of /newa/

Name	Last Modified	Size	DAP Response Links	Dataset Viewers
NEWA_MESOSCALE_ATLAS/	2019-07-03T10:37:46GMT	-	- - - - -	

[THREDDS Catalog XML](#)

Hyrax development sponsored by [NSF](#), [NASA](#), and [NOAA](#)

[OPeNDAP Hyrax \(1.16.0\) Documentation](#)

[Questions?](#) [Contact Support](#)


NORA3 & NEWA in details

- **Both** are derived from ERA5 reanalysis.
- **The New European Wind atlas (NEWA)** relies on the WRF model (no data assimilation) [1,2].
- **The NORwegian hindcast Archive (NORA3)** relies on HARMONIE-AROME (data assimilation of 2 m temperature and relative humidity [3]).

[1] Hahmann, A. N., Sīle, T., Witha, B., Davis, N. N., Dörenkämper, M., Ezber, Y., ... & Söderberg, S. (2020). The making of the new european wind atlas—part 1: Model sensitivity. *Geoscientific model development*, 13(10), 5053-5078.

[2] Dörenkämper, M., Olsen, B. T., Witha, B., Hahmann, A. N., Davis, N. N., Barcons, J., ... & Mann, J. (2020). The making of the new european wind atlas—part 2: Production and evaluation. *Geoscientific model development*, 13(10), 5079-5102.

[3] Haakenstad, H., Breivik, Ø., Furevik, B. R., Reistad, M., Bohlinger, P., & Aarnes, O. J. (2021). NORA3: A Nonhydrostatic High-Resolution Hindcast of the North Sea, the Norwegian Sea, and the Barents Sea. *Journal of Applied Meteorology and Climatology*, 60(10), 1443-1464.



Reference data from the offshore platform FINO1

Sensors selected:

Cup anemometer at 101 m

Wind vane at 91 m

Thermometer at 101 m

Year selected:

2009 (>8000 samples of 1h duration)



Metrics used to compare the datasets

Pearson correlation coefficient R^2 → Measures the linear correlation

Bias → Measures the systematic error


Root-mean square error (RMSE) → Measures the dispersion

They complement each other

Earth's mover distance (EMD) → Measures the distance between two probability density functions (pdfs)

Circular EMD (CEMD) → Same as EMD but for circular distributions

Mean wind speed comparison



Bias

NORA3: -0.11 ms^{-1}

NEWA: -0.31 ms^{-1}

Root-mean square error (RMSE)

NORA3: 1.3 ms^{-1}



NEWA: 2.0 ms^{-1}

Earth's mover distance (EMD)


NORA3: 0.13 ms^{-1}

NEWA: 0.26 ms^{-1}

Mean wind speed distributions



Mean wind direction distributions




Circular earth's mover distance (CEMD)

NORA3: 5.8°

NEWA: 7.0°

Larger-than-expected deviations due to calibration bias in the wind-vane data?

Sea surface temperature (SST)





MUR SST database collected from satellite data analysis [4]. Data available since 2002.

The buoy data were collected a few hundreds meters from the FINO1 platform (limited data availability).

[4] JPL MUR MEaSUREs Project. 2015. GHRSSST Level 4 MUR Global Foundation Sea Surface Temperature Analysis. Ver. 4.1. PO.DAAC, CA, USA.

Air temperature



Static stability (Brunt–Väisälä frequency, N)

$$N^2 = \frac{2 g [\theta(z_1) - \theta(z_2)]}{[\theta(z_1) + \theta(z_2)](z_1 - z_2)}$$

Here:

$$g = 9.81 \text{ ms}^{-2}$$


$$z_1 = 100 \text{ m}$$

$$z_2 = 0 \text{ m}$$

$\theta(z)$: potential temperature at height z

Improved agreement with measured data compared to a previous study [5]

[5] Barstad, I. (2016). Offshore validation of a 3 km ERA-Interim downscaling—WRF model's performance on static stability. *Wind Energy*, 19(3), 515-526.



Conclusions

Both NEWA and NORA3 show good performances to describe the distribution of mean wind speed and mean wind direction at FINO1 in 2009.

NORA3 slightly outperforms NEWA in terms of RMSE, bias, EMD, R^2 and CEMD

Both NEWA and NORA3 show encouraging results to characterize the static stability

Additional validation of the NORA3 database is ongoing.

See the poster [Offshore wind validation of the new NORA3 hindcast data with Lidar measurements](#), by J.M. Diezel (UiB)

Thank you

Acknowledgments:

The Federal Maritime and Hydrographic Agency of Germany (BSH) is acknowledged for providing the FINO1 data, available at <http://fino.bsh.de/>. We are grateful to Dr Andrea Hahmann and Dr Bjarke Tobias Olsen for their help regarding the implementation of the circular EMD function.



UNIVERSITETET I BERGEN